

An Overview of the Clinical and Molecular Genetics of Wilms' Tumor

Paul Grundy, MD, and Max Coppes, MD, PhD

INTRODUCTION

The treatment of Wilms' tumor is one of the remarkable success stories in pediatric oncology. Most children with Wilms' tumor in North America are registered on and subsequently treated according to National Wilms' Tumor Study Group (NWTSG) trials. In the past two decades, the NWTSG has contributed significantly to the development of therapeutic strategies for the management of these children. As a result, the majority of children with Wilms' tumor diagnosed in the 1990s are expected to survive and reach adulthood.

With a median age of 3.5 years at diagnosis, Wilms' tumor typically affects young children. However, the median age of diagnosis for certain children (those with bilateral Wilms' tumor, genitourinary anomalies, aniridia, or intralobar nephrogenic rests) is significantly younger, suggesting a predisposition for developing Wilms' tumor. Moreover, the associations of several genetically determined congenital anomalies (such as aniridia and Beckwith-Wiedemann syndrome [BWS]) with Wilms' tumor have also been of great help in identifying hereditary forms of Wilms' tumor and in determining possible locations of genes involved in its etiology.

A genetic model for the development of Wilms' tumor was originally proposed by Knudson and Strong in 1972 [1]. Based on analyses of age at diagnosis, and assuming constant mutation rates, they suggested that the development of Wilms' tumor required two rate-limiting genetic events. Children with genetic susceptibility would harbor an initial lesion in their germline, either inherited from a parent or resulting from a *de novo* mutation. The likelihood of a second genetic "hit" leading to tumorigenesis is thus greatly enhanced, compared with sporadic cases in which two rare independent somatic mutations are required. Subsequent genetic studies in a number of tumors have confirmed the Knudson model, demonstrating that the two postulated "genetic hits" constitute the inactivation of both alleles of a tumor suppressor gene. In many cases, the first allele is inactivated by a mutation within the gene itself, while the second allele is inactivated by a gross loss of chromosomal material, detected by so-called loss of heterozygosity (LOH) at one or more nearby polymorphic markers.

The two-event hypothesis predicts that susceptible individuals, such as familial cases, those with multifocal disease, and those with certain predisposing congenital

anomalies, have a lower median age at diagnosis than those without these characteristics. Studies by NWTSG investigators have confirmed the expected younger average age at diagnosis in children with multifocal or bilateral Wilms' tumor, although not for those with familial Wilms' tumor [2], suggesting that susceptibility to Wilms' tumor is complex and possibly involves more than one genetic locus and interactions between genetic and epigenetic factors.

It is now clear that several chromosomal loci are involved in the etiology of this childhood malignancy. These include one at chromosome 11p13 (*WT1*), one at chromosome 11p15 (*WT2*), possibly one at Xq25-27, and a yet unidentified "familial" locus. In addition, several loci are involved in Wilms' tumor progression, including one at chromosome 16q, one at chromosome 1p, and *p53* at chromosome 17p13.

CHROMOSOME 11p13 AND *WT1*

The description of children with the rare congenital WAGR syndrome, consisting of malformations including aniridia, genitourinary malformations, mental retardation, and Wilms' tumor [3] who had constitutional deletions of band p13 of chromosome 11 [4], was the first clue to the genomic location of a Wilms' tumor gene. The fact that this first event involved apparent chromosomal loss also suggested that the underlying gene might be a tumor suppressor, rather than an oncogene. The subsequent observation of tumor-specific LOH for 11p13 DNA markers in some sporadic tumors supported the two-hit hypothesis and further suggested that sporadic tumors involved the same locus as the WAGR cases [5-8].

This Wilms' tumor gene, *WT1*, has now been cloned [9-11] and shown to be a developmentally regulated [12] transcription factor of the zinc finger family [13] with expression restricted to the genitourinary system, spleen, dorsal mesentery of the intestines, muscles, central ner-

From the Departments of Pediatrics and Oncology, University of Alberta, and Department of Pediatrics, Cross Cancer Institute, Edmonton (P.G.), and Departments of Oncology and Pediatrics, University of Calgary, Tom Baker Cancer Center and Alberta Children's Hospital, Calgary (M.C.), Alberta, Canada.

Received July 13, 1995; accepted February 6, 1996.

Address reprint requests to Dr. Paul Grundy, Molecular Oncology Program, Cross Cancer Institute, 11560 University Ave., Edmonton, Alberta, Canada T6G 1Z2.

vous system, and mesothelium [14]. *WT1* is constitutionally deleted in all WAGR syndrome patients and there are examples of both bilateral [15] and sporadic [16] tumors with loss or independent mutations of the remaining allele, as predicted by the two-event tumor-suppressor model. Analyses in several series of sporadic Wilms' tumors, however, have found evidence for *WT1* mutation in only 5–10% of cases [17–19]. Thus, although *WT1* is undoubtedly a tumor-suppressor gene, it would appear to account for only a minority of Wilms' tumors.

Predisposition to Wilms' tumor is also a dominant feature of the rare Denys-Drash syndrome (DDS), which further consists of severe genitourinary anomalies (pseudohermaphroditism in males) and nephropathy progressing to renal failure (reviewed in [20]). Analysis of *WT1* in these patients has revealed the presence of constitutional, usually missense mutations, most frequently in the zinc finger region [21,22].

The majority of mutations described in Wilms' tumors are recessive, nonsense, or frame-shift mutations, whereas the majority of mutations described in DDS patients are dominant missense mutations [23]. All of the missense mutations described to date in DDS affect the DNA-binding ability of the predicted protein product and result in a far more severe phenotype than the one following complete constitutional inactivation of the *WT1* gene, i.e., WAGR syndrome.

CHROMOSOME 11p15 AND *WT2*

Wilms' tumor also occurs with increased frequency in BWS, a congenital and sometimes familial overgrowth and malformation condition variably consisting of macrosomia or hemihypertrophy, macroglossia, omphalocele, abdominal organomegaly, and ear pits or creases [24–26]. The BWS locus has been mapped to chromosome 11p15 based on constitutional karyotype abnormalities in rare patients [27], genetic linkage studies of families [28,29], and the rare occurrence of uniparental paternal isodisomy for 11p15 (the inheritance of two copies of one of the father's chromosomes 11 but neither of the mother's) [30,31].

In addition to the clinical association of Wilms' tumor and BWS, some sporadic Wilms' tumors have been identified with LOH restricted to markers at 11p15 not including 11p13 and *WT1* [28,32]. Intriguingly, an almost absolute bias to loss of the maternal allele has been found in Wilms' tumors [33], suggesting that *WT2*, the putative Wilms' tumor suppressor gene at 11p15, is subject to genomic imprinting [34]. That is, the gene is differentially marked, dependent on the parent of origin, such that the two alleles are not functionally equivalent [35]. Similarly, familial BWS segregates in a sex-specific fashion [28,36] and all reported cases of uniparental isodisomy have involved the paternal chromosome [30,31,37]. It is not yet

known, however, whether the BWS locus and *WT2* are the same or at contiguous loci. Evidence from in vitro growth suppression experiments does, however, suggest the existence of a tumor suppressor locus (loci) distinct from the BWS locus [38].

Two *WT2* candidate genes, *IGFII* and *H19*, have been studied. Both are imprinted in normal humans [39], although in the opposite direction. A possibly substantial fraction of Wilms' tumors (without LOH at the DNA level) has been found to have altered imprints with resultant overexpression of the growth factor *IGFII* [39,40] and loss of expression of the tumor-suppressor *H19* [39–42]. This has raised the possibility that only one genetic event might be required to effectively alter (enhance or inactivate) the relevant loci.

Besides BWS, Wilms' tumor also occurs with increased frequency in a very similar overgrowth syndrome, Simpson-Golabi-Behmel (SGB), which is transmitted in a sex-linked fashion and is linked to Xq25-27 [43,44]. The SGB locus has not been cloned and therefore has not been evaluated in sporadic tumors. It has been speculated that if the BWS locus represents an imprinted locus, that the SGB locus might represent the imprintor locus [44].

FAMILIAL WILMS' TUMOR

Familial Wilms' tumor, although rare, appears to segregate in an autosomal dominant fashion, albeit with low and possibly variable penetrance. The low incidence has complicated the search for a gene involved in familial Wilms' tumor. However, linkage analyses in three large families have excluded both 11p13 and 11p15 (i.e., *WT1* and *WT2*) from conferring susceptibility to Wilms' tumor [45,46]. A third locus, at 16q (see below), has also been excluded as the familial Wilms' tumor gene [47]. Consequently the location of the familial locus (loci) remains unknown.

WILMS' TUMOR PROGRESSION GENES

In addition to the three genetic loci discussed above (*WT1*, *WT2*, familial), which are implicated in the predisposition or genesis of Wilms' tumor, there is evidence for genetic loci which may be involved in the progression to more malignant or aggressive tumors. LOH at chromosome 16q has been observed in ~15–20% of Wilms' tumors [48–50], and at chromosome 1p in ~10% [50]. Constitutional alterations at these genomic locations have not been associated with a predisposition to Wilms' tumor, but tumor-specific loss of either region has been associated with an adverse outcome [50].

The tumor-suppressor gene, *p53*, also appears to play a role in a subset of tumors [51]. Tumor-specific mutations appear to be specifically associated with the so-called anaplastic histology, a subgroup with a particularly ad-

verse prognosis. Indeed, when individual foci of anaplastic cells have been dissected from the bulk of the tumor, the mutations have been confined to the anaplastic foci demonstrating that these genetic events take place within the already developed tumor [52].

However, *p53* mutations have also been documented in occasional favorable histology Wilms' tumors [53]. Nevertheless, the high correlation between *p53* mutations and anaplastic Wilms' tumors suggest that *p53* alterations are required for progression to the anaplastic phenotype, albeit with possible additional events.

Apart from *p53*, which is associated with cell cycle regulation and apoptosis, the nature of the other putative Wilms' progression genes is unknown.

CONCLUSIONS

The last few years have provided dramatic breakthroughs in understanding the genetic factors involved in Wilms' tumorigenesis and hence in normal kidney development. The implications of these findings for the clinical management of children with Wilms' tumor are only now becoming apparent.

Identification of the genetic defects involved in Wilms' tumor will allow more precise genetic counseling in the future. Children who present with sporadic aniridia and are known to be at risk of developing Wilms' tumor can now undergo molecular analysis of germline DNA. This analysis should readily distinguish children with a mutation restricted to the aniridia gene, *Pax6*, from those with a chromosomal deletion encompassing the neighboring *WT1* gene. Similarly, identification of the Wilms' tumor gene(s) residing at chromosome 11p15 and its relationship with the BWS gene may allow genetic counseling for children with hemihypertrophy, whose increased risk of developing embryonal malignancies is now poorly defined.

Faced with an unknown risk of metachronous bilateral Wilms' tumor, identification of children with germline mutations in the different Wilms' tumor genes may distinguish those at risk from the majority of sporadic unilateral cases.

Finally, once the genetic factors that underlie Wilms' tumorigenesis have been identified, analysis of parental exposures associated with the development of this tumor may prove of great value to evaluate the mechanisms that lead to the described genetic defects particularly with reference to genomic imprinting.

REFERENCES

- Knudson AG, Strong LC: Mutation and cancer: A model for Wilms' tumor of the kidney. *J Natl Cancer Inst* 48:313-324, 1972.
- Breslow N, Beckwith JB, Ciol M, Sharples K: Age distribution of Wilms' tumor: Report from the National Wilms' Tumor Study. *Cancer Res* 48:1653-1657, 1988.
- Miller RW, Fraumeni JF, Manning MD: Association of Wilms' tumor with aniridia, hemihypertrophy and other congenital anomalies. *N Engl J Med* 270:922-927, 1964.
- Riccardi VM, Sujansky E, Smith AC, Francke U: Chromosomal imbalance in the aniridia-Wilms' tumor association: 11p interstitial deletion. *Pediatrics* 61:604-610, 1978.
- Koufos A, Hansen MF, Lampkin BC, Workman ML, Copeland NG, Jenkins NA, Cawene WK: Loss of alleles at loci on human chromosome 11 during genesis of Wilms' tumour. *Nature* 309:170-172, 1984.
- Orkin SH, Goldman DS, Sallan SE: Development of homozygosity for chromosome 11p markers in Wilms' tumour. *Nature* 309:172-174, 1984.
- Reeve AE, Housiaux PJ, Gardner RJM, Chewings WE, Grindley RM, Milow LJ: Loss of a Harvey ras allele in sporadic Wilms' tumour. *Nature* 309:174-176, 1984.
- Fearon ER, Vogelstein B, Feinberg AP: Somatic deletion and duplication of genes on chromosome 11 in Wilms' tumours. *Nature* 309:176-178, 1984.
- Bonetta L, Kuehn SE, Huang A, Law DJ, Kalikin LM, Koi M, Reeve AE, Brownstein BH, Yeger H, Williams BRG, Feinberg AP: Wilms tumor locus on 11p13 defined by multiple CpG island-associated transcripts. *Science* 250:994-997, 1990.
- Gessler M, Poustka A, Cawene W, Neve RL, Orkin SH, Bruns GAP: Homozygous deletion in Wilms' tumours of a zinc-finger gene identified by chromosome jumping. *Nature* 343:774-778, 1990.
- Call KM, Glaser T, Ito CY, Buckler AJ, Pelletier J, Haber DA, Rose EA, Kral A, Yeger H, Lewis WH, Jones C, Housman DE: Isolation and characterization of a zinc finger polypeptide gene at the human chromosome 11 Wilms' tumor locus. *Cell* 60:509-520, 1990.
- Huang A, Campbell CE, Bonetta L, McAndrews-Hill M, Chilton-MacNeill S, Coppes MJ, Law DJ, Feinberg AP, Yeger H, Williams BRG: Tissue, developmental, and tumor-specific expression of divergent transcripts in Wilms' tumor. *Science* 250:991-994, 1990.
- Rauscher FJ, Morris JF, Tournay OE, Cook DM, Curran T: Binding of the Wilms' tumor locus zinc finger protein to the EGR-1 consensus sequence. *Science* 250:1259-1262, 1990.
- Pritchard-Jones K, Fleming S, Davidson D, Bickmore W, Porteous D, Gosden C, Bard J, Buckler A, Pelletier J, Housman D, Van Heyningen V, Hastie N: The candidate Wilms' tumour gene is involved in genitourinary development. *Nature* 346:194-197, 1990.
- Huff V, Miwa H, Haber D, Housman D, Strong LC, Saunders GF: Intragenic germinal deletion in bilateral Wilms' tumor: Evidence the WT33 identifies a WT gene (WT1) and support for the two-mutation model. *Am J Hum Genet* 48:997-1003, 1991.
- Lewis WH, Yeger H, Bonetta L, Chan HSL, Kang J, Junien C, Cowell J, Jones C, Dafoe LA: Homozygous deletion of a DNA marker from chromosome 11p13 in sporadic Wilms' tumor. *Genomics* 3:25-31, 1988.
- Coppes MJ, Liefers GJ, Paul P, Yeger H, Williams BRG: Homozygous somatic WT1 point mutations in sporadic unilateral Wilms' tumor. *Proc Natl Acad Sci USA* 90:1416-1419, 1993.
- Varanasi R, Bardeesy N, Ghahremani M, Petruzzi M-J, Nowak N, Adam MA, Grundy P, Shows TB, Pelletier J: Fine structure analysis of the *WT1* gene in sporadic Wilms' tumor. *Proc Natl Acad Sci USA* 91:3554-3558, 1994.
- Gessler M, Konig A, Arden K, Grundy P, Orkin S, Sallan S, Peters C, Ruyle S, Mandell J, Li F, Cawene W, Bruns G: Infrequent mutation of the *WT1* gene in 77 Wilms' tumors. *Hum Mutat* 3:212-222, 1994.
- Jadresic L, Leake J, Gordon I, Dillon MJ, Grant DB, Pritchard J, Risdon RA, Barratt TM: Clinicopathologic review of twelve children with nephropathy, Wilms tumor, and genital abnormalities (Drash syndrome). *J Pediatr* 117:717-725, 1990.

21. Huff V, Villalba F, Strong LC, Saunders G: Alteration of the WT1 gene in patients with Wilms' tumor and genitourinary anomalies. *Am J Hum Genet* 49:441, 1991.
22. Pelletier J, Bruening W, Cashtan CE, Mauer SM, Manivel JC, Striegel JE, Houghton DC, Junien C, Habib R, Fouser L, Fine RN, Silverman BL, Haber DA, Housman D: Germline mutations in the Wilms' tumor suppressor gene are associated with abnormal urogenital development in Denys-Drash syndrome. *Cell* 67:437–447, 1991.
23. Coppes MJ, Campbell EC, Williams BRG: "Wilms Tumor: Clinical and Molecular Characterization." New York: Springer-Verlag (in press).
24. Beckwith JB: Extreme cytomegaly of the adrenal fetal cortex, omphalocele, hyperplasia of kidneys and pancreas, and Leydig-cell hyperplasia: Another syndrome? Presented to the Western Society for Pediatric Research, Los Angeles, 1963.
25. Wiedemann HR: Complexe malformatif familial avec hernie ombilicale et macroglossie—un syndrome nouveau? *J Genet Hum* 13:223–232, 1964.
26. Beckwith JB: Macroglossia, omphalocele, adrenal cytomegaly, gigantism, and hyperplastic visceromegaly. *BD: OAS* 5:188–197, 1969.
27. Turleau C, De Grouchy J: Beckwith-Wiedemann syndrome clinical comparison between patients with and without 11p15 trisomy. *Ann Genet* 28:93–96, 1985.
28. Koufos A, Grundy P, Morgan K, Aleck KA, Hadro T, Lampkin BC, Kalbakji A, Cavenee WK: Familial Weidemann-Beckwith syndrome and a second Wilms' tumor locus both map to 11p15.5. *Am J Hum Genet* 44:711–719, 1989.
29. Ping AJ, Reeve AE, Law DJ, Young MR, Boehnke M, Feinberg AP: Genetic linkage of Beckwith-Weidmann syndrome to 11p15. *Am J Hum Genet* 44:720–723, 1989.
30. Henry I, Bonaiti-Pellie C, Chehensse V, Beldjord C, Schwartz C, Utermann G, Junien C: Uniparental paternal disomy in a genetic cancer-predisposing syndrome. *Nature* 351:665–667, 1991.
31. Grundy P, Telzerow P, Haber D, Berman B, Li F, Paterson MC, Garber J: Chromosome 11 uniparental isodisomy predisposing to embryonal neoplasms. *Lancet* 338:1079–1080, 1991.
32. Henry I, Grandjouan S, Couillin P, Barichard F, Huerre-Jeanpierre C, Glasser T, Philip T, Lenoir G, Chaussain JL, Junien C: Tumor-specific loss of 11p15.5 alleles in del11p13 Wilms' tumor and in familial adrenocortical carcinoma. *Proc Natl Acad Sci USA* 86:3247–3251, 1989.
33. Schroeder WT, Chao L, Dao DD, Strong LC, Pathak S, Riccardi V, Lewis WH, Saunders GF: Nonrandom loss of maternal chromosome 11 alleles in Wilms' tumors. *Am J Hum Genet* 40:413–420, 1987.
34. Wilkins RJ: Genomic imprinting and carcinogenesis. *Lancet* 1:329–331, 1988.
35. Scrabble HJ, Capienza C, Cavenee WK: Genetic and epigenetic losses of heterozygosity in cancer predisposition and progression. *Adv Cancer Res* 54:25–62, 1990.
36. Lubinsky M, Herrmann J, Kosseff AL, Opitz JM: Autosomal-dominant sex-dependent transmission of the Wiedemann-Beckwith syndrome. *Lancet* 1:932, 1974.
37. Henry I, Puech A, Riesewijk A, Ahnine L, Mannens M, Beldjord C, Bitoun P, Tourmade MF, Landrieu P, Junien C: Somatic mosaicism for partial paternal isodisomy in Wiedemann-Beckwith syndrome: A post-fertilization event. *Eur J Hum Genet* 1:19–29, 1993.
38. Koi M, Johnson LA, Kalikin LM, Little PFR, Nakamura Y, Feinberg AP: Tumor cell growth arrest caused by subchromosomal transferable DNA fragments from chromosome 11. *Science* 260:361–364, 1993.
39. Rainier S, Johnson LA, Dobry CJ, Ping AJ, Grundy PE, Feinberg AP: Relaxation of imprinted genes in human cancer. *Nature* 362:747–749, 1993.
40. Ogawa O, Eccles MR, Szeto J, McNoe LA, Yun K, Maw MA, Smith PJ, Reeve AE: Relaxation of insulin-like growth factor II gene imprinting implicated in Wilms' tumour. *Nature* 362:749–751, 1993.
41. Steenman MJC, Rainier S, Dobry CJ, Grundy P, Horon IL, Feinberg AP: Loss of imprinting of IGF2 is linked to reduced expression and abnormal methylation of H19 in Wilms' tumour. *Nature Genet* 7:433–439, 1994.
42. Moulton T, Crenshaw T, Hao Y, Moosikasuwan J, Lin N, Dembitzer F, Hensle T, Weiss L, McMorrow L, Loew T, Kraus W, Gerald W, Tycko B: Epigenetic lesions at the H19 locus in Wilms' tumour patients. *Nature Genet* 7:440–447, 1994.
43. Xuan JY, Besner A, Ireland M, Hughes-Benzie RM, MacKenzie AE: Mapping of Simpson-Golabi-Beckmel syndrome to Xq25-q27. *Hum Mol Genet* 3:133–138, 1994.
44. Hughes-Benzie RM, Hunter AGW, Allanson JE, MacKenzie AE: Simpson-Golabi-Beckmel syndrome associated with renal dysplasia and embryonal tumor: Localization of the gene to Xqcen-q21. *Am J Med Genet* 43:428–435, 1992.
45. Grundy P, Koufos A, Morgan K, Li FP, Meadows AT, Cavenee WK: Familial predisposition to Wilms' tumour does not map to the short arm of chromosome 11. *Nature* 336:374–376, 1988.
46. Huff V, Compton DA, Chao L, Strong LC, Geiser CF, Saunders GF: Lack of linkage of familial Wilms' tumour to chromosomal band 11p13. *Nature* 336:377–378, 1988.
47. Huff V, Reeve AE, Leppert M, Strong LC, Douglass EC, Geiser CF, Li FP, Meadows A, Callen DF, Lenoir G, Saunders GF: Non-linkage of 16q markers to familial predisposition to Wilms' tumor. *Cancer Res* 52:6117–6120, 1992.
48. Maw MA, Grundy PE, Millow LJ, Eccles MR, Dunn RS, Smith PJ, Feinberg AP, Law DJ, Paterson MC, Telzerow PE, Callen DF, Thompson AD, Richards RI, Reeve AE: A third Wilms' tumor locus on chromosome 16q. *Cancer Res* 52:3094–3098, 1992.
49. Coppes MJ, Bonetta L, Huang A, Hoban P, Chilton-MacNeill S, Campbell CE, Weksberg R, Yeger H, Reeve AE, Williams BRG: Loss of heterozygosity mapping in Wilms' tumor indicates the involvement of three distinct regions and a limited role for nondisjunction or mitotic recombination. *Genes Chromosomes Cancer* 5:326–334, 1992.
50. Grundy PE, Telzerow PE, Breslow N, Moksness J, Huff V, Paterson MC: Loss of heterozygosity for chromosomes 16q and 1p in Wilms' tumors predicts an adverse outcome. *Cancer Res* 54:2331–2333, 1994.
51. Bardeesy N, Falkoff D, Petruzzi M-J, Nowak N, Zabel B, Adam M, Aguiar MC, Grundy P, Shows T, Pelletier J: Anaplastic Wilms' tumour, a subtype displaying poor prognosis, harbours p53 gene mutations. *Nature Genet* 7:91–97, 1994.
52. Bardeesy N, Beckwith JB, Pelletier J: Clonal expansion and attenuated apoptosis in Wilms' tumor are associated with p53 gene mutations. *Cancer Res* 55:215–219, 1995.
53. Malkin D, Sexsmith E, Yeger H, Williams BRG, Coppes MJ: Mutations of the p53 tumor suppressor gene occur infrequently in Wilms' tumor. *Cancer Res* 54:2077–2079, 1994.